arXiv:1412.4971v1 [math.CA] 16 Dec 2014

Rényi entropy and Tsallis entropy associated with positive linear operators

Ioan Raşa*

Subjclass: 41A36, 94A17, 33C45
Keywords: Rényi entropy, Tsallis entropy, positive linear operators, Legendre polynomials.

Abstract

This article is a continuation of my paper [arxiv: 1409.1015v2]. Rényi and Tsallis entropies are associated to positive linear operators and properties of some functions related to these entropies are investigated.

1 Introduction

This paper is a continuation of [9]. In that article we considered discrete positive linear operators of the form

$$Lf(x)=\sum_k f(x_k)a_k(x), \quad a_k(x)\geq 0, \quad \sum_k a_k(x)=1$$

for x in some interval $I\subset\mathbb{R}$. In investigating the degree of non-multiplicativity of L an important role was played by the function $S(x)=\sum_k a_k^2(x)$; see [6].

On the other hand, for each fixed $x\in I$ the numbers $(a_k(x))_k$ form a probability distribution. In this context $-\log S(x)$ is a Rényi entropy [10] and $1-S(x)$ is a Tsallis entropy [11]. So the properties of the function $S(x)$ are relevant also in the study of these entropies.

Some properties of $S(x)$ were investigated in [5]-[9]. In Section 2 we continue to study such properties in the case of discrete operators. Section 3 is devoted to some multivariate operators; see also [1]. In the last section we consider integral operators of the form $Lf(x)=\int_I K(x,t)f(t)dt$ with $K(x,t)\geq 0$, $\int_I K(x,t)dt=1$, $x\in I$. Recall that in this case the associated Rényi entropy is defined by $-\log\int_I K^2(x,t)dt$, and the Tsallis entropy by $1-\int_I K^2(x,t)dt$.

*Department of Mathematics, Technical University of Cluj-Napoca, Memorandumului Street 28, 400114 Cluj-Napoca, Romania, ioan.rasa@math.utcluj.ro

2 A conjecture from [9] and some of its consequences

We shall use the notation from [9]. In particular, we consider the function $S_{n,c}$ defined on the interval I_c. Conjecture 7.1 in [9] reads as follows:

$$(C) \quad \log S_{n,c} \text{ is a convex function.}$$

Let us examine some consequences of this conjecture. With $X := x(1+cx)$ and $X' = 1+2cx$, (3.10) in [9] can be written as

$$XX'S''_{n,c}(x) + (4(n+c)X+1)S'_{n,c}(x) + 2nX'S_{n,c}(x) = 0. \tag{2.1}$$

(C) is equivalent to $S''_{n,c}S_{n,c} \geq (S'_{n,c})^2$, and due to (2.1) both of them are equivalent to

$$XX'\left(\frac{S'_{n,c}}{S_{n,c}}\right)^2 + (4(n+c)X+1)\frac{S'_{n,c}}{S_{n,c}} + 2nX' \leq 0.$$

This leads immediately to

Theorem 2.1 *Conjecture (C) is equivalent to*

$$(C') \quad \frac{S'_{n,c}(x)}{S_{n,c}(x)} \text{ is between } z_1(x) \text{ and } z_2(x),$$

where

$$z_1(x) = \frac{-\sqrt{(1+4cX)^2+(4nX)^2} - (1+4cX) - 4nX}{2XX'},$$

$$z_2(x) = \frac{\sqrt{(1+4cX)^2+(4nX)^2} - (1+4cX) - 4nX}{2XX'}.$$

Let us consider the case $c = 0$. Then the function $K_n(x) := S_{n,0}(x)$ is defined for $x \in [0, +\infty)$. In [9, (6.4)] it was proved that

$$K_n(x) \leq \frac{1}{\sqrt{4nx+1}}, \quad x \geq 0. \tag{2.2}$$

Recall also that I_0 is the modified Bessel function of first kind of order zero, and (see [9, (3.7)])

$$I_0(x) = e^x K_n\left(\frac{x}{2n}\right), \quad x \geq 0. \tag{2.3}$$

Combining (2.2) and (2.3) we get

$$I_0(x) \leq \frac{\exp x}{\sqrt{2x+1}}, \quad x \geq 0. \tag{2.4}$$

Corollary 2.2 *Under the hypothesis that* $\log S_{n,0}$ *is convex, we have*

$$\frac{-\sqrt{1+(4nt)^2}-1-4nt}{2t} \le \frac{K_n'(t)}{K_n(t)} \le \frac{\sqrt{1+(4nt)^2}-1-4nt}{2t}, \quad t>0, \quad (2.5)$$

$$K_n^2(x) \le \frac{2\exp\left(\sqrt{1+(4nx)^2}-1-4nx\right)}{\sqrt{1+(4nx)^2}+1}, \quad x \ge 0, \quad (2.6)$$

$$I_0^2(x) \le \frac{2\exp\left(\sqrt{1+4x^2}-1\right)}{\sqrt{1+4x^2}+1}, \quad x \ge 0. \quad (2.7)$$

Proof. (2.5) is a direct consequence of Theorem 2.1. (2.6) can be obtained from the second inequality in (2.5) by integrating with respect to t between 0 and x. (2.7) follows from (2.6) and (2.3).

In order to compare (2.2) with (2.6), and (2.4) with (2.7), it is easy to check that

$$\frac{2\exp\left(\sqrt{1+(4nx)^2}-1-4nx\right)}{\sqrt{1+(4nx)^2}+1} \le \frac{1}{4nx+1}, \quad x \ge 0,$$

$$\frac{2\exp\left(\sqrt{1+4x^2}-1\right)}{\sqrt{1+4x^2}+1} \le \frac{\exp 2x}{2x+1}, \quad x \ge 0.$$

Now let us consider the case $c=-1$. Then the function $F_n(x) := S_{n,-1}(x)$ is defined for $x \in [0,1]$. Let $x \in [0,1/2)$ and $t = \frac{2x^2-2x+1}{1-2x}$. Then (see also [9, Section 4]) $t \in [1,+\infty)$, $x = \frac{1-t+\sqrt{t^2-1}}{2}$, $X = x(1-x) = \sqrt{t^2-1}(t-\sqrt{t^2-1})/2$, $X' = 1-2x = t-\sqrt{2-1}$, $\frac{dt}{dx} = \frac{4X}{1-4X}$, $t = (1-2X)/X'$, $\sqrt{t^2-1} = 2X/X'$.

Let $(P_n(t))_{n\ge 0}$ be the Legendre polynomials. Then (see [7], [8], [9, (4.6)])

$$F_n(x) = \left(t-\sqrt{t^2-1}\right)^n P_n(t). \quad (2.8)$$

This yields

$$\frac{F_n'(x)}{F_n(x)} = \left(\frac{P_n'(t)}{P_n(t)} - \frac{n}{\sqrt{t^2-1}}\right)\frac{dt}{dx},$$

and consequently

$$\frac{P_n'(t)}{P_n(t)} - \frac{n}{\sqrt{t^2-1}} = \frac{1-4X}{4X}\frac{F_n'(x)}{F_n(x)}. \quad (2.9)$$

Corollary 2.3 *Under the hypothesis that* $\log S_{n,-1}$ *is convex, we have*

$$\frac{F_n'(x)}{F_n(x)} \le \frac{\sqrt{(1-4X)^2+(4nX)^2}-(1-4X)-4nX}{2XX'}, \quad x \in \left[0,\frac{1}{2}\right), \quad (2.10)$$

$$\frac{P_n'(t)}{P_n(t)} \le \frac{\sqrt{4n^2(t^2-1)+(t-\sqrt{t^2-1})^2}-(t-\sqrt{t^2-1})}{2(t^2-1)}, \quad t>1. \quad (2.11)$$

Proof. (2.10) is a consequence of Theorem 2.1. (2.11) follows from (2.10) and (2.9).

The following inequality was proved in [8, (1.2)]:

$$\frac{n(n+1)}{2t+(n-1)\sqrt{t^2-1}} \leq \frac{P_n'(t)}{P_n(t)}, \quad t \geq 1. \tag{2.12}$$

Using it and (2.9), we get

$$-\frac{2nX'}{1+(n-3)X} \leq \frac{F_n'(x)}{F_n(x)}, \quad x \in \left[0, \frac{1}{2}\right]. \tag{2.13}$$

Other lower and upper bounds for $\frac{P_n'(t)}{P_n(t)}$ can be found in [8]. In particular, from [8, Theorems 2 and 3] we have

$$\frac{P_n'(t)}{P_n(t)} \leq \frac{2n^2}{t+(2n-1)\sqrt{t^2-1}}, \quad t \geq 1, \tag{2.14}$$

$$\frac{P_n'(t)}{P_n(t)} \leq \frac{n^2(2n+1)}{(n+1)t+(2n^2-1)\sqrt{t^2-1}}, \quad t \geq 1. \tag{2.15}$$

(2.11) and (2.14) can be compared and we get

$$\frac{\sqrt{4n^2(t^2-1)+(t-\sqrt{t^2-1})^2}-(t-\sqrt{t^2-1})}{2(t^2-1)} \leq \frac{2n^2}{t+(2n-1)\sqrt{t^2-1}}, \quad t>1.$$

The inequality

$$\frac{\sqrt{4n^2(t^2-1)+(t-\sqrt{t^2-1})^2}-(t-\sqrt{t^2-1})}{2(t^2-1)} \leq \frac{n^2(2n+1)}{(n+1)t+(2n^2-1)\sqrt{t^2-1}}, \quad t>1,$$

is equivalent to

$$\frac{t}{t+\sqrt{t^2-1}} \geq \frac{3n+2}{4n+3}.$$

This last inequality is true for t approaching 1, and false for t approaching $+\infty$.

Let us remark that (2.15) yields by integration

$$P_n(t) \leq (t+\sqrt{t^2-1})^{\frac{n(2n^2-1)}{2n^2-n-2}} \left(t+\frac{2n^2-1}{n+1}\sqrt{t^2-1}\right)^{-\frac{n(n+1)}{2n^2-n-2}}, \quad t \geq 1. \tag{2.16}$$

This inequality is stronger than

$$P_n(t) \leq (t+\sqrt{t^2-1})^{\frac{n(2n-1)}{2(n-1)}} \left(t+(2n-1)\sqrt{t^2-1}\right)^{-\frac{n}{2(n-1)}}, \quad t \geq 1, n \geq 2, \tag{2.17}$$

which can be obtained from (2.14).

We conclude this section with a remark concerning the function $A_{n,c} := \frac{S'_{n,c}}{S_{n,c}}$. By using [9, (3.10)], or (2.1), we deduce easily that $A_{n,c}$ satisfies the Riccati equation

$$x(1+cx)(1+2cx)(A'_{n,c}+A^2_{n,c})+(4(n+c)x(1+cx)+1)A_{n,c}+2n(1+2cx)=0.$$

3 Multivariate operators

First, consider the classical Bernstein operators on the canonical simplex of $\mathbb{R}^2$. The sum of the squared fundamental Bernstein polynomials is in this case

$$\begin{aligned} R_n(x,y) &:= \sum_{i+j\le n}\left(\frac{n!}{i!j!(n-i-j)!}\right)^2 x^{2i}y^{2j}(1-x-y)^{2(n-i-j)} \\ &= \sum_{j=0}^{n}\sum_{i=0}^{n-j}\binom{n}{j}^2\binom{n-j}{i}^2 x^{2i}y^{2j}(1-x-y)^{2(n-i-j)}, \end{aligned}$$

for $x\ge 0$, $y\ge 0$, $x+y\le 1$; see [2, (6.3.6)], [3, Sect. 3.1.2].

Let $y\in[0,1)$ be fixed. Then for $x\in[0,1-y]$ we have

$$R_n(x,y)=\sum_{j=0}^{n}\binom{n}{j}^2 y^{2j}(1-y)^{2(n-j)}\sum_{i=0}^{n-j}\binom{n-j}{i}^2\left(\frac{x}{1-y}\right)^{2i}\left(1-\frac{x}{1-y}\right)^{2(n-j-i)}.$$

For each $j\in\{0,1,...,n\}$,

$$\sum_{i=0}^{n-j}\binom{n-j}{i}^2\left(\frac{x}{1-y}\right)^{2i}\left(1-\frac{x}{1-y}\right)^{2(n-j-i)} = F_{n-j}\left(\frac{x}{1-y}\right),$$

where $F_{n-j}=S_{n-j,-1}$. It is known (see [5], [6], [8], [9]) that F_{n-j} is convex on $[0,1]$. It follows that for each fixed $y\in[0,1)$, the function $R_n(\cdot,y)$ is convex on $[0,1-y]$. In other words, R_n is convex on each segment parallel to Ox. Similarly we see that R_n is convex on each segment parallel to a side of the canonical triangle of $\mathbb{R}^2$. This means that R_n is *axially-convex*; concerning this terminology see [2, p. 407], [3, Sect. 3.5].

Now consider the classical Bernstein operators on the square $[0,1]^2$: see [2, (6.3.101)], [3, Sect. 3.1.5]. The sum of the squared fundamental Bernstein polynomials is in this case

$$Q_n(x,y)=\sum_{i=0}^{n}\sum_{j=0}^{n}\binom{n}{i}^2\binom{n}{j}^2 x^{2i}(1-x)^{2n-2i}y^{2j}(1-y)^{2n-2j}=F_n(x)F_n(y).$$

It is easy to verify that the following three statements are equivalent:

i) $\log F_n$ is convex on $[0,1]$;

ii) Q_n is convex on $[0,1]^2$;

iii) $\log Q_n$ is convex on $[0,1]^2$.

4 Entropy and variance. Integral operators

Let I be an interval and L a positive linear operator on a space of functions defined on I, containing the functions $e_i(x) = x^i$, $i = 0, 1, 2$. Suppose that $Le_0 = e_0$.

The *variance* associated with L is the function

$$V(x) := Le_2(x) - (Le_1(x))^2, \quad x \in I.$$

If L is a discrete operator of the form $Lf(x) = \sum_k f(x_k)a_k(x)$, let $S(x) := \sum_k a_k^2(x)$, $x \in I$. If L is an integral operator of the form $Lf(x) = \int_I K(x,t)f(t)dt$, let $S(x) := \int_I K^2(x,t)dt$, $x \in I$.

In both cases the Rényi entropy associated with L is $-\log S(x)$, and the Tsallis entropy is $1 - S(x)$, $x \in I$.

Example 4.1

Let $L_{n,c}f(x) = \sum_{j=0}^{\infty} f\left(\frac{j}{n}\right) p_{n,j}^{[c]}(x)$, see [9, Sect. 2]. Then $S(x) = S_{n,c}(x)$ and $V(x) = V_{n,c}(x) = \frac{x(1+cx)}{n}$. According to [9, (3.5), (3.8)],

$$S_{n,c}(x) = \frac{1}{\pi}\int_0^{\pi} \left(1 + 4ncV_{n,c}(x)\sin^2\frac{\varphi}{2}\right)^{-n/c} d\varphi, \quad c \neq 0, \tag{4.1}$$

$$S_{n,0}(x) = \frac{1}{\pi}\int_0^{\pi} \exp\left(-4n^2V_{n,0}(x)\sin^2\frac{\varphi}{2}\right)d\varphi. \tag{4.2}$$

Example 4.2

For the Kantorovich operators [2, p. 333] we have $S_n(x) = (n+1)F_n(x)$ and $V_n(x) = (n+1)^{-2}\left(nx(1-x) + \frac{1}{12}\right)$.

Example 4.3

Consider the Gauss-Weierstrass operators [2, p. 310], [4, p. 114]:

$$W_rf(x) = \int_{\mathbb{R}} (4\pi r)^{-1/2} \exp\left(-\frac{(t-x)^2}{4r}\right) f(t)dt, \quad r > 0.$$

Then $V_r(x) = 2r$ and $S_r(x) = (8\pi r)^{-1/2}$, $x \in \mathbb{R}$.

Generally speaking, for a convolution operator

$$Lf(x) = \int_{\mathbb{R}} \varphi(x-t)f(t)dt \text{ we have } V(x) = \int_{\mathbb{R}} s^2\varphi(s)ds - \left(\int_{\mathbb{R}} s\varphi(s)ds\right)^2$$

and $S(x) = \int_{\mathbb{R}} \varphi^2(s)ds$, so that V and S are constant functions.

Example 4.4

For the Post-Widder operators [4, p. 114],

$$V_n(x) = \frac{x^2}{n} \text{ and } S_n(x) = \binom{2n-2}{n-1} 2^{1-2n} \frac{n}{x}, \quad x > 0.$$

Example 4.5

Consider the Durrmeyer operators [2, p. 335].

In this case

$$V_n(x) = \frac{n+1}{(n+2)^2(n+3)}(2nx(1-x)+1),$$

$$S_n(x) = \sum_{k=0}^{2n} c_{n,k} \binom{2n}{k} x^k (1-x)^{2n-k},$$

where

$$c_{n,k} := \frac{(n+1)^2}{2n+1} \binom{2n}{k}^{-2} \sum_{j=0}^{k} \binom{n}{j}^2 \binom{n}{k-j}^2, \quad k = 0, 1, ..., 2n,$$

where, as usual, $\binom{n}{m} = 0$ if $m > n$.

It is easy to see that $c_{n,2n-k} = c_{n,k}$, $k = 0, 1, ..., 2n$.

Conjecture 4.6 *The sequence $(c_{n,k})_{k=0,1,..,2n}$ is convex and, consequently, the function S_n is convex on $[0, 1]$.*

Example 4.7

For the genuine Bernstein-Durrmeyer operators, defined by

$$U_n f(x) = f(0)b_{n,0}(x) + f(1)b_{n,n}(x) + (n-1)\sum_{k=1}^{n-1} b_{n,k}(x) \int_0^1 b_{n-2,k-1}(t) f(t) dt,$$

with $b_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$, we have

$$V_n(x) = \frac{2x(1-x)}{n+1}$$

and

$$S_n(x) = (1-x)^{2n} + x^{2n} +$$
$$+ \frac{(n-1)^2}{2n-3} \sum_{k,j=1}^{n-1} \binom{n-2}{k-1} \binom{n-2}{j-1} \binom{n}{k} \binom{n}{j} \binom{2n-4}{k+j-2}^{-1} x^{k+j} (1-x)^{2n-k-j}.$$

Remark 4.8 *In Examples 4.1-4.4, and also in Example 4.5 under Conjecture 4.6, the functions $V(x)$, $1 - S(x)$ and $-\log S(x)$ are all increasing or all decreasing on suitable subintervals of I. In other words, the variance, the Tsallis entropy and the Rényi entropy are synchronous functions.*

Acknowledgement

The author is grateful to Dr. Gabriela Raluca Mocanu for inspiring discussions.

References

[1] A. Acu and M.-D. Rusu, New results concerning Chebyshev-Grüss-type inequalities via discrete oscillations, Appl. Math. Comput. 243 (2014), 585-593.

[2] F. Altomare and M. Campiti, Korovkin-type Approximation Theory and its Applications, de Gruyter, Berlin-New York, 1994.

[3] F. Altomare, M. Cappelletti Montano, V. Leonessa, I. Raşa, Markov Operators, Positive Semigroups and Approximation Processes, de Gruyter Studies in Mathematics, vol. 61, Walter de Gruyter, Berlin, 2014.

[4] Z. Ditzian and V. Totik, Moduli of Smoothness, Springer-Verlag, New York, 1987.

[5] I. Gavrea and M. Ivan, On a conjecture concerning the sum of the squared Bernstein polynomials, Appl. Math. Comput. 241 (2014), 70-74.

[6] H. Gonska, I. Raşa and M.-D. Rusu, Chebyshev-Grüss-type inequalities via discrete oscillations, Bul. Acad. Stiinte Repub. Mold. Mat. no. 1 (74) (2014), 63-89. arxiv 1401.7908 [math.CA].

[7] T. Neuschel, Unpublished manuscript, Univ. of Trier, 2pp., 2012.

[8] G. Nikolov, Inequalities for ultraspherical polynomials. Proof of a conjecture of I. Raşa, J. Math. Anal. Appl. 418 (2014), 852-860.

[9] I. Raşa, Special functions associated with positive linear operators, arxiv: 1409.1015v2 [math.CA].

[10] A. Rényi, On measures of entropy and information, Proc. Fourth Berkeley Symp. Math. Statist. Prob., Vol. 1, Univ. of California Press 1961, 547-561.

[11] C. Tsallis, Possible generalization of Boltzmann-Gibbs statistics, J. Stat. Phys. 52 (1988), 479-487.